

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07K 5/06, A61K 37/02</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/20527</b> <b>(43) International Publication Date:</b> 15 September 1994 (15.09.94)
<b>(21) International Application Number:</b> PCT/US94/02706 <b>(22) International Filing Date:</b> 11 March 1994 (11.03.94)  <b>(30) Priority Data:</b> 08/030,770 12 March 1993 (12.03.93) US 08/031,187 12 March 1993 (12.03.93) US  <b>(60) Parent Application or Grant</b> <b>(63) Related by Continuation</b> US 08/031,287 (CIP) Filed on 12 March 1993 (12.03.93)  <b>(71) Applicant (for all designated States except US):</b> ARRIS PHARMACEUTICAL CORPORATION [US/US]; 385 Oyster Point Boulevard, Suite 3, South San Francisco, CA 94080 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SPEAR, Kerry [US/US]; 5732 Colton Boulevard, Oakland, CA 94611 (US). JOHN-SON, Charles [US/US]; 2670 Hilgard Avenue, Berkeley, CA 94709 (US). GSCHWEND, Heinz, W. [US/US]; 20192 Osprey Drive, Bodega Bay, CA 94923 (US).		<b>(74) Agent:</b> DOW, Karen, B.; Townsend and Townsend Kourie and Crew, Stuart Street Tower, 20th floor, One Market Plaza, San Francisco, CA 94105 (US).  <b>(81) Designated States:</b> AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> COMPOSITIONS AND METHODS FOR THE TREATMENT OF IMMUNOMEDIATED INFLAMMATORY DISORDERS  <b>(57) Abstract</b>  Compositions and methods for the prevention and treatment of immunomediated inflammatory disorders, especially for those disorders associated with the respiratory tract, are provided. More particularly, a tryptase inhibitor, typically a hydroxyaroyl or hydroxyheteroaroyl substituted dipeptide, is administered. Also provided by this invention are pharmaceutical compositions, typically aerosol or topical, as well as aerosol devices for administering these compositions intranasally.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5                   **COMPOSITIONS AND METHODS FOR THE TREATMENT OF**  
                    **IMMUNOMEDIATED INFLAMMATORY DISORDERS**

**CROSS-REFERENCE TO RELATED APPLICATION**

                    This application is a continuation-in-part of  
                    copending application Serial No. 08/031,187, which is related  
10                   to the subject matter of U.S. Patent Application Serial No.  
                    08/030,770, both applications filed on March 12, 1993, titled  
                    "COMPOSITIONS AND METHODS FOR THE TREATMENT OF IMMUNOMEDIATED  
                    INFLAMMATORY DISORDERS" and "METHODS FOR THE TREATMENT OF  
                    HYPERRESPONSIVENESS ASSOCIATED WITH CHRONIC ASTHMA",  
15                   respectively. These copending applications are expressly  
                    incorporated herein by reference.

**BACKGROUND OF THE INVENTION**

20                   1.    Field of the Invention

                    This invention relates generally to compositions and  
                    methods for the prevention and treatment of immunomediated  
                    inflammatory disorders. More particularly, the invention  
                    relates to the prevention and treatment of inflammatory  
                    diseases associated with the respiratory tract, such as asthma  
25                   and allergic rhinitis. The compositions and methods of the  
                    present invention are especially useful for preventing or  
                    treating the late phase bronchoconstriction and airway  
                    hyperresponsiveness associated with chronic asthma.

30                   2.    Description of the Background Art

                    Asthma is a complex disease involving multiple  
                    biochemical mediators for both its acute and chronic  
                    manifestations. Asthma frequently is characterized by  
                    progressive development of hyperresponsiveness of the trachea  
35                   and bronchi to both immunospecific allergens and generalized  
                    chemical or physical stimuli. The hyperresponsiveness of  
                    asthmatic bronchiolar tissue is believed to result from chronic  
                    inflammation reactions, which irritate and damage the  
                    epithelium lining the airway wall and promote pathological  
40                   thickening of the underlying tissue. Bronchial biopsy studies

have indicated that even patients with mild asthma have features of inflammation in the airway wall.

One initiator of the inflammatory sequence is an allergic response to inhaled allergens. Leukocytes carrying  
5 IgE receptors, notably mast cells and basophils, but also including monocytes, macrophages, and eosinophils, are present in the epithelium and underlying smooth muscle tissues of bronchi, where they are activated initially by binding of specific inhaled antigens to the IgE receptors. Activated mast  
10 cells release a number of preformed or primary chemical mediators of the inflammatory response and enzymes. Furthermore, numerous secondary mediators of inflammation are generated *in situ* by enzymatic reactions of activated mast cells, including superoxide and lipid derived mediators. In  
15 addition, several large molecules are released by degranulation of mast cells: proteoglycans, peroxidase, arylsulfatase B, and notably the proteases tryptase and chymotryptic proteinase (chymase). See "Drug Therapy of Asthma", Chap. 62, 1054-54.

This chemical release from mast cells probably  
20 accounts for the early bronchiolar constrictor response that occurs in susceptible individuals after exposure to airborne allergens. The early asthmatic reaction is maximal at around fifteen minutes after allergen exposure; recovery occurs over the ensuing one to two hours. In 25-35% of individuals, the  
25 early asthmatic reaction is followed by a further decline in respiratory function which begins within a few hours and is maximal between six and twelve hours post-exposure. This late asthmatic reaction is accompanied by a marked increase in the number of inflammatory cells infiltrating bronchiolar smooth  
30 muscle and epithelial tissues, and spilling into the airways. These cells include eosinophils, neutrophils, and lymphocytes, all of which are attracted to the site by release of mast cell derived chemotactic agents. The infiltrating cells themselves become activated during the late reaction phase. The late  
35 asthmatic response is believed to be a secondary inflammatory reaction mediated in part by the secretory activity of macrophages.

A related set of inflammatory reactions occurs in the upper respiratory tract mucosa, usually in response to airborne allergens. As in asthma, mast cells are activated by crosslinking of IgE molecules to particular antigens. In allergic, perennial or vasomotor rhinitis, mast cells may be activated in the absence of discernible exposure to a particular antigen. In either case, activated mast cells release primary and secondary mediators of inflammation upon degranulation. Eosinophils and macrophages are attracted to the site to perpetuate the inflammation reaction. Nasal epithelial tissue destruction often occurs in late-phase reactions.

Tryptase is the major secretory protease of human mast cells and is proposed to be involved in neuropeptide processing and tissue inflammation. Mature human tryptase is a glycosylated, heparin-associated tetramer of heterogenous, catalytically active subunits. The tryptase monomer's amino acid sequence, like its gene structure, has no close counterpart among the numerous other serine proteinases that have been characterized. See, e.g., Vanderslice et al. (1990) Proc. Natl. Acad. Sci. USA 87:3811-3815; Miller et al. (1990) J. Clin. Invest. 86:864-870; Miller et al. (1989) J. Clin. Invest. 84:1188-1195; and Vanderslice et al. (1989) Biochemistry 28:4148-4155.

Tryptase is stored in mast cell secretory granules. After mast cell activation, human tryptase can be measured readily in a variety of biologic fluids. For example, after anaphylaxis, tryptase appears in the bloodstream, where it remains detectable for several hours. See Schwartz et al. (1987) N. Engl. J. Med. 316:1622-1626. Its appearance has been detected in samples of nasal and lung lavage fluid from atopic subjects challenged with specific antigen. See Castells and Schwartz 1988) J. Allerg. Clin. Immunol. 82:348-355 and Wenzel et al. (1988) Am. Rev. Resp. Dis. 141:563-568. Tryptase levels in lung lavage fluid obtained from atopic asthmatics increase after endobronchial allergen challenge. *Id.* Some smokers of cigarettes have striking elevations of bronchoalveolar lavage fluid tryptase levels compared to nonsmoking controls, a

finding that provides some support for the hypothesis that release of proteinases from activated mast cells could contribute to lung destruction in smoker's emphysema. See Kalenderian et al. (1988) Chest 94:119-123. In addition, tryptase has been shown to be a potent mitogen for fibroblasts, suggesting its involvement in pulmonary fibrosis and interstitial lung diseases. See Ruoss et al. (1991) J. Clin. Invest. 88:493-499.

Tryptase has been implicated in a variety of biological processes, including degradation of vasodilating and bronchorelaxing neuropeptides (see Caughey et al. (1988) J. Pharmacol. Exp. Ther. 244:133-137; Franconi et al. (1988) J. Pharmacol. Exp. Ther. 248:947-951; and Tam et al. (1990) Am. J. Respir. Cell Mol. Biol. 3:27-32) and modulation of bronchial responsiveness to histamine (see Sekizawa et al. (1989) J. Clin. Invest. 83:175-179). These studies suggest that tryptase possibly increases bronchoconstriction in asthma by destroying bronchodilating peptides.


Additionally, tryptase has been shown to cleave fibrinogen  $\alpha$ -chains, as well as high molecular weight kininogen with a possible release of kinins and thus, may play a role with heparin as a local anticoagulant. The ability of tryptase to activate prostromelysin (pro-MMP-3) and procollagenase (pro-MMP-1) via MMP-3 suggests that tryptase also may be involved in tissue inflammation and remodeling. This finding also intimates that tryptase may play a role in joint destruction in rheumatoid arthritis. In addition, tryptase has been shown to cleave calcitonin gene-related peptide. As this peptide is implicated in neurogenic inflammation, tryptase could be a factor in the regulation of flare reaction in cutaneous neurogenic inflammation. See Caughey (1991) Am. J. Respir. Cell Mol. Biol. 4:387-394.

Asthma has become the most common chronic disease in industrialized countries. To date, conventional methods and therapeutic agents have not proved to be effective in the treatment of asthma or other immunomediated inflammatory disorders. For these reasons it would be desirable to provide improved compositions and methods which avoid the disadvantages

f these conventional agents and methods while providing effective treatment for these diseases.

#### SUMMARY OF THE INVENTION

The present invention provides novel tryptase inhibitors comprising compounds of Formula I:



or a pharmaceutically acceptable salt thereof, wherein:


Ar is hydroxyl substituted aryl or hydroxyl substituted heteroaryl, wherein the hydroxyl is positioned ortho to the amide side chain and wherein if Ar is hydroxyl substituted aryl, the aromatic ring bearing the amide side chain is not substituted with halogen and does not bear a lower alkyl group on the position ortho to the hydroxyl;

R<sup>1</sup> is hydrogen, lower alkyl, arylalkyl, or heteroarylalkyl;


R<sup>2</sup> is hydrogen or lower alkyl;

R<sup>3</sup> is selected from the group consisting of:


6




5




10




15




20




25



30



and







wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; r is an integer from 0-5; s is an integer from 0-2; t is an integer from 1-3; u is 1 or 2; v is an integer from 3-6; and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is  $-\text{NH}-$  or  $-\text{CH}_2-$ ;



$R^4$  is lower alkyl, substituted arylalkyl or substituted heteroarylalkyl and  $R^5$  and  $R^6$  are independently selected from the group consisting of hydrogen, lower alkyl, substituted arylalkyl, and substituted heteroarylalkyl; or  $R^4$  and  $R^5$  together with the nitrogen and carbon to which they are attached, form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $R^6$  is hydrogen; or  $R^4$  and  $R^6$  together with the nitrogen and carbon to which they are attached, form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $R^5$  is hydrogen; and

$R^7$  is  $-OR^8$  or  $-NR^8R^9$ , wherein  $R^8$  and  $R^9$  are independently selected from the group consisting of hydrogen, lower alkyl, aryl, arylalkyl, or heteroarylalkyl or  $R^8$  and  $R^9$  together with the nitrogen to which they are attached, form a substituted 5-membered or 6-membered heterocycle.


In a preferred embodiment, Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;  $R^1$  is hydrogen;  $R^2$  is hydrogen;  $R^3$  is selected from the group consisting of:




wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is  $-\text{NH}-$  or  $-\text{CH}_2-$ ;

either  $\text{R}^4$  is lower alkyl and  $\text{R}^5$  and  $\text{R}^6$  are hydrogen; or  $\text{R}^4$  and  $\text{R}^5$ , together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $\text{R}^6$  is hydrogen; and  $\text{R}^7$  is  $-\text{OH}$ ,  $-\text{OCH}_3$ ,  $-\text{NH}_2$ , 3'-aminocarboxy-1'-piperidyl, or  $-\text{N}(\text{CH}_3)_2$ .

A particularly preferred tryptase inhibitor of Formula I is Compound 3 of Tables I and II:



Another preferred tryptase inhibitor of Formula I is Compound 15 of Tables I and II:



The compounds described herein are useful for the prevention and treatment of immunomediated inflammatory disorders, and particularly those associated with the

respiratory tract, including asthma, and particularly the hyperresponsiveness phase associated with chronic asthma, and allergic rhinitis. Thus, the present invention also provides a method for treating immunomediated inflammatory disorders wherein a patient having an immunomediated inflammatory disorder that is susceptible to treatment with a tryptase inhibitor receives, or is administered, a therapeutically effective dose or amount of a compound of the present invention.

The invention also provides for pharmaceutical compositions of the compounds described herein. These pharmaceutical compositions can be in a variety of forms including oral dosage forms, as well as injectable and infusible solutions. Typically, when used for the treatment or prevention of the asthma, and particularly the hyperresponsiveness associated with chronic asthma, these pharmaceutical compositions are in an aerosol form of powders or solutions. When used for the treatment of immunomediated inflammatory skin conditions, the compounds of the instant invention are used in combination with a non-toxic, pharmaceutically acceptable topical carrier. The compounds of the instant invention can be used in combination with antiinflammatories or other asthma therapies, such as  $\beta$ -adrenergic agonists, antiinflammatory corticosteroids, anticholinergics, and the like.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph showing specific lung resistance in sheep as a function of time in hours post-antigen challenge. Open squares indicate control values, and filled circles indicate values for the same animal after administration of Compound 3 of Tables I and II.

Figure 2 is a graph showing production of hyperresponsiveness of sheep to carbachol-induced bronchoconstriction at 24 hours post-allergen challenge when Compound 3 of Tables I and II was administered before carbachol. The dark, solid bar corresponds to the control, baseline. The lighter, solid bar corresponds to the drug,

baseline. The hatched bar corresponds to control, post-antigen. The white bar corresponds to the drug, post-antigen.

#### DESCRIPTION OF SPECIFIC EMBODIMENTS

##### 5 I. Definitions and General Parameters

The following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

"Immunomediated inflammatory disorder" includes  
10 generally those diseases associated with mast cell mediator release and susceptible to treatment with a tryptase inhibitor. Examples of such disorders include diseases of immediate type hypersensitivity such as asthma, allergic rhinitis, urticaria and angioedema, and eczematous dermatitis (atopic dermatitis),  
15 and anaphylaxis, as well as hyperproliferative skin disease, peptic ulcers, inflammatory bowel disorder, inflammatory skin conditions, and the like.

"Hyperresponsiveness" refers to late phase bronchoconstriction and airway hyperreactivity associated with  
20 chronic asthma. Hyperresponsiveness of asthmatic bronchiolar tissue is believed to result from chronic inflammation reactions, which irritate and damage the epithelium lining the airway wall and promote pathological thickening of the underlying tissue.

25 "Halogen" refers to fluorine, bromine, chlorine, and iodine atoms.

"Hydroxyl" refers to the group -OH.

"Lower alkyl" refers to a cyclic, branched or straight chain alkyl group of one to six carbon atoms. This  
30 term is further exemplified by such groups as methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, i-butyl (or 2-methylpropyl), cyclopropylmethyl, i-amyl, n-amyl, and hexyl.

"Aryl" or "Ar" refers to an aromatic carbocyclic group having a single ring (e.g., phenyl), multiple rings  
35 (e.g., biphenyl) or multiple condensed rings in which at least one ring is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl, anthryl, or phenanthryl), which can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl,

lower alkoxy, lower alkylthio, trifluoromethyl, lower acyloxy, aryl, heteroaryl and hydroxy. However, according to the instant invention, the aromatic ring bearing the amide side chain cannot be further substituted with halogen. In addition, 5 the aromatic ring bearing the amide side chain cannot possess a lower alkyl group ortho to the hydroxyl group (i.e., meta to the amide side chain).

"Heterocycle" refers to a saturated, unsaturated, or aromatic carbocyclic group having a single ring (e.g., 10 morpholino, pyridyl or furyl) or multiple condensed rings (e.g., naphthyridinyl, quinoxalyl, quinolinyl, indolizinyl or benzo[b]thienyl) and having at least one hetero atom, such as N, O or S, within the ring, which can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl, 15 lower alkoxy, lower alkylthio, trifluoromethyl, lower acyloxy, and hydroxy. The term "heteroaryl" or "HetAr" refers to a heterocycle in which at least one heterocyclic ring is aromatic.

"Arylalkyl" refers to the group -R-Ar where Ar is an 20 aryl group and R is straight-chain or branched-chain aliphatic group. Arylalkyl groups can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, lower acyloxy, and hydroxy.

"Heteroarylalkyl" refers to the group -R-HetAr where 25 HetAr is an heteroaryl group and R is straight-chain or branched-chain aliphatic group. Heteroarylalkyl groups can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, lower acyloxy, and hydroxy.

30 "Pharmaceutically acceptable salt" refers to those salts which retain the biological effectiveness and properties of the parent compound and which are not biologically or otherwise undesirable.

"Pharmaceutically or therapeutically acceptable 35 carrier" refers to a carrier medium which does not interfere with the effectiveness of the biological activity of the active ingredients and which is not toxic to the host or patient.

"Stereoisomer" refers to a chemical compound having the same molecular weight, chemical composition, and constitution as another, but with the atoms grouped differently. That is, certain identical chemical moieties are at different orientations in space and, therefore, when pure, has the ability to rotate the plane of polarized light. However, some pure stereoisomers may have an optical rotation that is so slight that it is undetectable with present instrumentation. The compounds of the instant invention may have one or more asymmetrical carbon atoms and therefore include various stereoisomers. All stereoisomers are included within the scope of the invention.


"Treatment" or "treating" refers to any administration of a tryptase inhibitor *in vitro* or *in vivo* and includes:

- (i) inhibiting the symptoms of the disease;
- (ii) lessening or inhibiting the long term effects of the disease; and/or
- (iii) relieving the symptoms of the disease.

## II. Tryptase Inhibitors

The present invention provides compositions comprising an effective serine protease inhibitor, and more particularly a tryptase inhibitor, that is useful for reducing immunomediated-inflammatory disorders, and particularly bronchoconstriction induced by allergenic challenge in an asthmatic animal.

Tryptase inhibitors are substances which slow down or prevent tryptase activity. According to one aspect of the present invention, the tryptase inhibitors will comprise a compound of Formula I:



or a pharmaceutically acceptable salt thereof,  
wherein:

Ar is hydroxyl substituted aryl or hydroxyl  
substituted heteroaryl, wherein the hydroxyl is positioned  
5 ortho to the amide side chain and wherein if Ar is hydroxyl  
substituted aryl, the aromatic ring bearing the amide side  
chain is not substituted with halogen and does not bear a lower  
alkyl group on the position ortho to the hydroxyl;

10  $R^1$  is hydrogen, lower alkyl, arylalkyl, or  
heteroarylalkyl;

$R^2$  is hydrogen or lower alkyl;

$R^3$  is selected from the group consisting of:

15


20

25


30

35


14




5




10





15




20



25



30







wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; r is an integer from 0-5; s is an integer from 0-2; t is an integer from 1-3; u is 1 or 2; v is an integer from 3-6; and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is  $-\text{NH}-$  or  $-\text{CH}_2-$ ;



$R^4$  is lower alkyl, substituted arylalkyl, or substituted heteroarylalkyl and  $R^5$ , and  $R^6$  are independently selected from the group consisting of hydrogen, lower alkyl, substituted arylalkyl, and substituted heteroarylalkyl; or  $R^4$  and  $R^5$  together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $R^6$  is hydrogen; or  $R^4$  and  $R^6$  together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $R^5$  is hydrogen; and

$R^7$  is  $-OR^8$  or  $-NR^8R^9$ , wherein  $R^8$  and  $R^9$  are independently selected from the group consisting of hydrogen, lower alkyl, aryl, arylalkyl, or heteroarylalkyl or  $R^8$  and  $R^9$  together with the nitrogen to which they are attached, form a substituted 5-membered or 6-membered heterocycle.


In a preferred embodiment, Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;  $R^1$  is hydrogen;  $R^2$  is hydrogen;  $R^3$  is selected from the group consisting of:




wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is  $-\text{NH}-$  or  $-\text{CH}_2-$ ;

either  $\text{R}^4$  is lower alkyl and  $\text{R}^5$  and  $\text{R}^6$  are hydrogen; or  $\text{R}^4$  and  $\text{R}^5$  together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $\text{R}^6$  is hydrogen; and  $\text{R}^7$  is  $-\text{OH}$ ,  $-\text{OCH}_3$ ,  $-\text{NH}_2$ , 3'-aminocarboxy-1'-piperidyl, or  $-\text{N}(\text{CH}_3)_2$ .


A preferred tryptase inhibitor is Compound 3 of Tables I and II:



An additional preferred tryptase inhibitor is Compound 15 of Tables I and II:



Another preferred tryptase inhibitor is Compound 21 of Tables I and II:



The tryptase inhibitors of the present invention can  
5 be obtained by known techniques from readily available starting  
materials, as described in greater detail below.

Compounds of this invention can, depending on the  
nature of the functional groups, form addition salts with  
various inorganic and organic acids and bases. These salts can  
10 be formed with inorganic acids such as hydrochloric acid,  
hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid  
and the like, and organic acids such as acetic acid, propionic  
acid, glycolic acid, pyruvic acid, oxalic acid, malic acid,  
malonic acid, succinic acid, maleic acid, fumaric acid,  
15 tartaric acid, citric acid, benzoic acid, cinnamic acid,  
mandelic acid, methanesulfonic acid, ethanesulfonic acid,  
p-toluenesulfonic acid, salicylic acid and the like.

Salts can also be formed from a carboxylic acid  
residue via treatment with alkali metals or alkali metal bases,  
20 such as alkali metal hydroxides and alkali metal alkoxides, or  
alkaline earth metals or alkaline earth metal bases, such as  
alkaline earth metal hydroxides and alkaline earth metal  
alkoxides. In addition, salts can be formed from a carboxylic  
acid and an organic base, such as trimethylamine, diethylamine,  
25 ethanolamine, piperidine, isopropylamine, choline, caffeine,  
and the like.

The salts can be formed by conventional means, as by  
reacting the free acid or base forms of the product with one or

more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water, which is then removed *in vacuo* or by freeze-drying or by exchanging the cations of an existing salt for another  
5 cation on a suitable ion exchange resin.

### III. In Vitro and In Vivo Testing

*In vitro* protocols for screening potential inhibitors as to their ability to inhibit tryptase are known in the art.  
10 See, e.g., Sturzebecher et al. (1992) Biol. Chem. Hoppe-Seyler 373:1025-1030. Typically, these assays measure the tryptase-induced hydrolysis of peptide-based chromogenic substances. Details of an exemplary procedure are described below.

In addition, the activity of the compounds of the  
15 present invention can be evaluated *in vivo* in one of the numerous animal models of asthma. See Larson, "Experimental Models of Reversible Airway Obstruction", in The Lung: Scientific Foundations, Crystal, West et al., eds., Raven Press, New York, 1991; Warner et al. (1990) Am. Rev. Respir.  
20 Dis. 141:253-257. An ideal animal model would duplicate the chief clinical and physiological features of human asthma, including: airway hyperresponsiveness to chemical mediators and physical stimuli; reversal of airway obstruction by drugs useful in human asthma ( $\beta$ -adrenergics, methylxanthines,  
25 corticosteroids, and the like); airway inflammation with infiltration of activated leukocytes; and chronic inflammatory degenerative changes, such as basement membrane thickening, smooth muscle hypertrophy, and epithelial damage. Species used as animal models include mice, rats, guinea pigs, rabbits,  
30 dogs, and sheep. All have some limitations, and the proper choice of animal model depends upon the question which is to be addressed.

The initial asthmatic response can be evaluated in guinea pigs, and dogs, and particularly, with a basenji-  
35 greyhound cross strain which develops nonspecific airway hyperresponsiveness to numerous nonallergenic substances, such as methacholine and citric acid. Certain selected sheep exhibit a dual response after antigen challenge with *Ascaris*

proteins. In dual responding animals, the initial asthmatic response (IAR) is followed by a late asthmatic response (LAR) at 6-8 hours post-exposure. Hypersensitivity to the cholinergic agonist carbachol increases at 24 hours after antigen challenge in those animals which exhibit LAR.

The allergic sheep model was used to evaluate the potential antiasthmatic effects of the compounds of the present invention. Administration of compositions comprising aerosolized solutions of the compounds of the instant invention to allergic sheep prior to or following exposure to specific allergens demonstrates that such compositions substantially lessen or abolish the late asthmatic response and consequent hyperresponsiveness.

The compounds of this invention are also useful for the treatment of other immunomediated inflammatory disorders in which tryptase activity contributes to the pathological condition. Such diseases include inflammatory diseases associated with mast cells, such as rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, inflammatory bowel disease, peptic ulcer and various skin conditions.

The efficacy of the compounds of the instant invention for the treatment of the vast majority of immunomediated inflammatory disorders can be evaluated by either *in vitro* or *in vivo* procedures. Thus, the anti-inflammatory efficacy of the compounds of the instant invention can be demonstrated by assays well known in the art, for example, the Reversed Passive Arthus Reaction (RPAR)-PAW technique (see, e.g., Ganguly et al. (1992) U.S. Patent No. 5,126,352). Assays for determining the therapeutic value of compounds in the treatment of various skin conditions, such as hyperproliferative skin disease, are well known in the art, for example, the Arachidonic Acid Mouse Ear Test (*id*). The compounds of the instant invention can be evaluated for their antiulcer activity according to the procedures described in Chiu et al. (1984) Archives Internationales de Pharmacodynamie et de Therapie 270:128-140.

#### IV. In Vivo Administration

According to this invention, a therapeutically or pharmaceutically effective amount of a tryptase inhibitor and particularly, a compound of Formula I, is administered to a patient suffering from an immunomediated inflammatory disorder. According to one embodiment, the compositions of the present invention are useful for preventing or ameliorating asthma. In using the compositions of the present invention in a treatment of asthma, the compounds may be administered prophylactically prior to exposure to allergen or other precipitating factor, or after such exposure. The compounds of the instant invention are particularly useful in ameliorating the late-phase tissue destruction seen in both seasonal and perennial rhinitis. Another aspect of the present invention is directed to the prevention and treatment of other immunomediated inflammatory disorders associated with mast cells such as urticaria and angioedema, and eczematous dermatitis (atopic dermatitis), and anaphylaxis, as well as hyperproliferative skin disease, peptic ulcers, and the like.

The compositions containing the compounds can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, compositions are administered to a patient already suffering from a disease, as described above, in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as "therapeutically effective amount or dose." Amounts effective for this use will depend on the severity and course of the disease, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician.

In prophylactic applications, compositions containing the compounds of the invention are administered to a patient susceptible to or otherwise at risk of a particular disease. Such an amount is defined to be a "prophylactically effective amount or dose." In this use, the precise amounts again depend on the patient's state of health, weight, and the like.

Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary.

Subsequently, the dosage or the frequency of administration, or both, can be reduced, as a function of the symptoms, to a level at which the improved condition is retained. When the symptoms have been alleviated to the desired level, treatment can cease.

5 Patients can, however, require intermittent treatment on a long-term basis upon any recurrence of the disease symptoms.

In general, a suitable effective dose of the tryptase inhibitor will be in the range of 0.1 to 1000 milligram (mg) per recipient per day, preferably in the range of 1 to 100 mg  
10 per day. The desired dosage is preferably presented in one, two, three, four or more subdoses administered at appropriate intervals throughout the day. These subdoses can be administered as unit dosage forms, for example, containing 5 to 1000 mg, preferably 10 to 100 mg of active ingredient per unit  
15 dosage form.

The composition used in these therapies can be in a variety of forms. These include, for example, solid, semi-solid and liquid dosage forms, such as tablets, pills, powders, liquid solutions or suspensions, liposomes, injectable and  
20 infusible solutions. The preferred form depends on the intended mode of administration and therapeutic application.

While it is possible to administer the active ingredient of this invention alone, it is preferable to present it as part of a pharmaceutical formulation. The formulations  
25 of the present invention comprise at least one compound or inhibitor of this invention in a therapeutically or pharmaceutically effective dose together with one or more pharmaceutically or therapeutically acceptable carriers and optionally other therapeutic ingredients. Various  
30 considerations are described, e.g., in Gilman et al. (eds) (1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's supra. Methods for administration are discussed therein, e.g., for oral, intravenous, intraperitoneal, or intramuscular  
35 administration, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, NJ.

Typically, when the compounds of the instant invention are to be used in the treatment of asthma or allergic rhinitis, they will be formulated as aerosols. The term "aerosol" includes any gas-borne suspended phase of the compounds of the instant invention which is capable of being inhaled into the bronchioles or nasal passages. Specifically, aerosol includes a gas-borne suspension of droplets of the compounds of the instant invention, as may be produced in a metered dose inhaler or nebulizer, or in a mist sprayer. Aerosol also includes a dry powder composition of a compound of the instant invention suspended in air or other carrier gas, which may be delivered by insufflation from an inhaler device, for example.

For solutions used in making aerosols of the present invention, the preferred range of concentration of the compounds of the instant invention is 0.1-100 milligrams (mg)/milliliter (mL), more preferably 0.1-30 mg/mL, and most preferably, 1-10 mg/mL. Usually the solutions are buffered with a physiologically compatible buffer such as phosphate or bicarbonate. The usual pH range is 5 to 9, preferably 6.5 to 7.8, and more preferably 7.0 to 7.6. Typically, sodium chloride is added to adjust the osmolarity to the physiological range, preferably within 10% of isotonic. Formulation of such solutions for creating aerosol inhalants is discussed in Remington's Pharmaceutical Sciences, see also, Ganderton and Jones, Drug Delivery to the Respiratory Tract, Ellis Horwood (1987); Gonda (1990) Critical Reviews in Therapeutic Drug Carrier Systems 6:273-313; and Raeburn et al. (1992) J. Pharmacol. Toxicol. Methods 27:143-159.

Solutions of the compounds of the instant invention may be converted into aerosols by any of the known means routinely used for making aerosol inhalant pharmaceuticals. In general, such methods comprise pressurizing or providing a means of pressurizing a container of the solution, usually with an inert carrier gas, and passing the pressurized gas through a small orifice, thereby pulling droplets of the solution into the mouth and trachea of the animal to which the drug is to be administered. Typically, a mouthpiece is fitted to the outlet



of the orifice to facilitate delivery into the mouth and trachea.

In one embodiment, devices of the present invention comprise solutions of the compounds of the instant invention  
5 connected to or contained within any of the conventional means for creating aerosols in asthma medication, such as metered dose inhalers, jet nebulizers, or ultrasonic nebulizers. Optionally such device may include a mouthpiece fitted around the orifice.

10 In an embodiment for the treatment of allergic rhinitis, a device may comprise a solution of a compound of the instant invention in a nasal sprayer.

A dry powder comprising a compound of the instant invention, optionally with an excipient, is another embodiment  
15 of the present invention. This may be administered by a drug powder inhaler containing the above described powder.

It should, of course, be understood that the methods of this invention can be used in combination with other agents for the treatment of immunomediated inflammatory disorders, and  
20 particularly asthma.  $\beta$ -Adrenergic agonists are especially useful in these combinations, because they provide symptomatic relief of the initial asthmatic response, whereas the compounds of the present invention provide relief for the late asthmatic response. Preferred  $\beta$ -adrenergic agonists in these solutions  
25 include any of the usual  $\beta$ -agonists employed for the relief of asthma, such as albuterol, terbutaline, formoterol, fanoterol, or prenaline.

Other agents useful in combination with the compounds of the instant invention include anticholinergics, such as  
30 ipratropium bromide, and antiinflammatory corticosteroids (adrenocortical steroids) such as beclomethasone, triamcinolone, flurisolide, or dexamethasone.

The compounds of the inventions can also be used in the treatment of immunomediated inflammatory skin conditions,  
35 such as urticaria and angioedema, eczematous dermatitis, and hyperproliferative skin disease, e.g., psoriasis, in mammals. As a result of the topical administration of a compound of Formula I, a remission of the symptoms can be expected. Thus,

one affected by an immunomediated inflammatory skin condition can expect a decrease in scaling, erythema, size of the plaques, pruritus, and other symptoms associated with the skin condition. The dosage of medicament and the length of time  
5 required for successfully treating each individual patient may vary, but those skilled in the art will be able to recognize these variations and adjust the course of therapy accordingly.

Also included within the invention are preparations for topical application to the skin comprising a compound of  
10 Formula I, typically in concentrations in the range of from about 0.001% to 10%, together with a non-toxic, pharmaceutically acceptable topical carrier. These topical preparations can be prepared by combining an active ingredient according to this invention with conventional pharmaceutical  
15 diluents and carriers commonly used in topical dry, liquid, cream and aerosol formulations. Ointment and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Such bases may include water and/or an oil such as liquid paraffin  
20 or a vegetable oil such as peanut oil or castor oil. Thickening agents which may be used according to the nature of the base include soft paraffin, aluminum stearate, cetostearyl alcohol, propylene glycol, polyethylene glycols, woolfat, hydrogenated lanolin, beeswax, and the like.

25 Lotions may be formulated with an aqueous or oily base and will, in general, also include one or more of the following: stabilizing agents, emulsifying agents, dispersing agents, suspending agents, thickening agents, coloring agents, perfumes, and the like.

30 Powders may be formed with the aid of any suitable powder base, e.g., talc, lactose, starch, and the like. Drops may be formulated with an aqueous base or non-aqueous base also comprising one or more dispersing agents, suspending agents, solubilizing agents, and the like.

35 The topical pharmaceutical compositions according to this invention may also include one or more preservatives or bacteriostatic agents, e.g., methyl hydroxybenzoate, propyl hydroxybenzoate, chlorocresol, benzalkonium chlorides, and the

like. The topical pharmaceutical compositions also can contain other active ingredients such as antimicrobial agents, particularly antibiotics, anesthetics, analgesics, and antipruritic agents.

5           The compounds of this invention are also useful in the treatment of peptic ulcers. More specifically, they display chemotherapeutic activity which enables them to relieve the symptoms of peptic ulcer disease and stress ulceration, and to promote the healing of gastric and/or duodenal ulcers. The  
10       compounds can be used in conjunction with other therapeutic agents, such as anti-inflammatory and/or analgesic agents, such as aspirin, indomethacin, phenylbutazone, ibuprofen, naproxen, tolemtim, and the like.

          The pharmaceutical compositions can be administered  
15       by parenteral or oral administration for prophylactic and/or therapeutic treatment. The pharmaceutical compositions can be administered in a variety of unit dosage forms depending upon the method of administration. For example, unit dosage forms suitable for oral administration include powder, tablets,  
20       pills, capsules and dragees.

          The pharmaceutical compositions can be administered intravenously. Thus, this invention provides compositions for intravenous administration which comprise a solution of the compound dissolved or suspended in an acceptable carrier,  
25       preferably an aqueous carrier. A variety of aqueous carriers can be used, e.g., water, buffered water, 0.4% saline, and the like. These compositions will sometimes be sterilized by conventional, well known sterilization techniques, or can be sterile filtered. The resulting aqueous solutions can be  
30       packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous solution prior to administration. The compositions can contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and  
35       buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, and the like.

For solid compositions, conventional nontoxic solid carriers can be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable nontoxic composition is formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 0.1-95% of active ingredient, preferably about 20%.

In order that the invention described herein can be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only, and are not to be construed as limiting this invention in any manner.

## EXPERIMENTAL

### I. General

Starting materials not described herein are available commercially, are known, or can be prepared by methods known in the art. Compounds of Formula I can be prepared by techniques known in the art using either solid phase or solution phase synthesis techniques. Starting materials for the preparation of compounds of Formula I are known in the art or can be prepared by methods well known to those skilled in the art. For example, techniques for solid phase synthesis of polypeptides are described, for example, in Solid Phase Peptide Synthesis: A Practical Approach, (eds. E. Atherton and R. C. Sheppard) IRL Press at Oxford University Press (1989) and in Merrifield, J. Amer. Chem. Soc. 85:2149-2156 (1963). Peptide coupling chemistry is also described in The Peptides, Vol. 1, (eds. Gross, E., and J. Meienhofer), Academic Press, Orlando (1979). Other techniques include those of Geysen et al., J. Imm. Meth., (1987) 102:259-274; Houghten et al., Nature (1991) 354:84-86.

In the process described herein for the preparation of compounds of this invention, the requirements for protecting groups are generally well recognized by one skilled in the art

of organic chemistry. Accordingly, the use of appropriate protecting groups is necessarily implied by the processes contained herein, although not expressly illustrated.

Isolation and purification of the compounds and intermediates described herein can be effected, if desired, by any suitable separation or purification procedure such as, for example, filtration, extraction, crystallization, column chromatography, thin-layer chromatography or thick-layer chromatography, high-pressure liquid chromatography, or a combination of these procedures. Specific illustrations of suitable separation and isolation procedures can be had by reference to the examples hereinbelow. However, other equivalent separation or isolation procedures can, of course, also be used.

The Sakaguchi assay was for the detection of arginine and peptides containing arginine. See Stewart and Young "Solid Phase Peptide Synthesis", 2d Ed., Pierce Chemical Company, p. 114. Solution I containing 0.01%  $\alpha$ -naphthol and 5% urea in 95% ethanol was prepared. Solution II was prepared by dissolving 2 gram (g) bromine in 100 milliliters (mL) of 8% aqueous sodium hydroxide. To Solution I was added 5 sodium hydroxide pellets. The sample to be analyzed was spotted on a thin layer chromatography (TLC) plate. The TLC plate was then sprayed with Solution I. The TLC plate was dried in air and then sprayed with solution II. A red spot indicated the existence of an aminoimino methane group.

## II. Solid Phase Synthesis of Compounds of Formula I

### A. Preparation of Compound 3 of Tables I and II

4-(2',4'-Dimethoxyphenyl-fluorenylmethyloxycarbonyl (Fmoc)-aminomethyl)-phenoxy resin (Rink resin; Bachem, CA; 3.5 grams (g) at a loading of 0.289 milliequivalent/gram (meq/g), 1.01 millimoles (mmol)) was suspended in 30 milliliters (mL) of a 1:1 solution (volume/volume (v/v)) of N,N-dimethylformamide (DMF):toluene containing 30% (by volume) piperidine. The reaction mixture was agitated for 5 minutes and was then filtered. Additional piperidine solution (30 mL) was added and the mixture was agitated for an additional 5 minutes. After

filtration, the resin beads were washed sequentially with DMF (6x) and methylene chloride (6x).

Fmoc-L-proline (Milligen; 6.06 mmol), benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate (BOP, Novabiochem; 6.06 mmol), and 1-hydroxybenzotriazole (HOBT; Aldrich; 6.06 mmol) were dissolved in ~30 mL of DMF. The resulting clear solution was added to the Rink resin and the slurry was agitated (with nitrogen bubbling) for 4 hours. The reaction mixture was filtered and the resin beads were washed as described above. The Fmoc group was removed with piperidine as described above.

A solution of Fmoc-L-arginine(PMC) (Milligen; 6.06 mmol), BOP (6.06 mmol), and HOBT (6.06 mmol) in DMF (30 mL) was added to the resin beads and the slurry was agitated for 4 hours and was then filtered, washed, treated with piperidine, and washed again as described above.

A solution of 1-hydroxy-2-naphthoic acid (Aldrich; 1.14 gm, 6.06 mmol), 1,3-diisopropylcarbodiimide (DIPCDI; Aldrich; 0.949 mL, 6.06 mmol) and HOBT (0.819 g, 6.06 mL) in DMF (45 mL) was added and the reaction mixture was agitated (mechanical rocker arm) for 7 hours. After filtration, washing and piperidine treatment (described above), the resin was washed sequentially with DMF (6x), methylene chloride (6x) and methanol (6x). The resin was suspended in methanol and agitated (mechanical rocker arm) overnight (15 hours). The resin was filtered and washed twice with methanol.

The white, free flowing solid was then suspended in a solution of trifluoroacetic acid (TFA):anisole:water (90:5:5, 30 mL). The resin immediately turned pink and then darkened over the course of 1-2 minutes to a deep red color. After agitating for 5 minutes the reaction mixture was filtered. This procedure was repeated 2 additional times with fresh cleavage reagent, except that contact time was increased to 10 minutes for each iteration.

The TFA solutions were combined and the resulting clear, orange solution was allowed to sit at room temperature for 3.5 hours. Concentration *in vacuo* afforded a pale yellow oil which was kept *in vacuo* (<1 torr) for 3 hours. The oil was

dissolved in 50% aqueous acetonitrile (100 mL) and lyophilized to afford the crude product as an off-white solid (471 mg).

Analytical HPLC (Polymer Labs 100A PLRP column, 1.0 x 150 millimeters (mm), eluting with a linear gradient of 20 to 45% acetonitrile over 13 minutes at a flow of 0.1 mL/min) showed that the crude product was essentially a single compound (retention time 10.35 minutes, >98% by UV absorbance). Purification was accomplished via HPLC using a C<sub>18</sub> silica gel column (Vydac; 22 x 250 mm, 15-20  $\mu$ m, 300 A, eluting with a linear gradient of 20 to 35% over 30 minutes at a flow of 10 mL/minute). Multiple injections of 30-35 mg were required. Like fractions (retention time ~ 44-48 minutes) were combined and lyophilized to afford a fluffy white solid (351 mg) which was a single compound by HPLC.

Electrospray mass spectroscopy (calculated molecular weight = 440.49, found  $M^{+1}$  = 441.1) and NMR spectroscopy (proton and <sup>13</sup>C) are consistent with the expected structure. <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  7.89 (d, J=8Hz, 1 H), 7.78 (d, J=6Hz, 1 H), 7.76 (d, J=6Hz, 1 H), 7.57 (dt, J=8, 1Hz, 1 H), 7.48 (dt, J=8, 1Hz, 1 H), 7.29 (d, J=8Hz, 1 H), ~4.95 (obscured, 1 H), 4.49 (dd, J=8, 5Hz, 1 H), 3.97 (dt, J=10, 7Hz, 1 H), 3.73 (dt, J=10, 7Hz, 1 H), 3.23 (t, J=7Hz, 2 H), 2.29 (m, 1 H), 1.7-2.2 (m, 8 H).

B. Preparation of Other Compounds of Formula I

By following the procedure of Part A above and substituting for 1-hydroxy-2-naphthoic acid the following compounds:

- 2-hydroxy-4-methylbenzoic acid;
- 3-hydroxy-2-quinoxalinecarboxylic acid;
- 3-hydroxy-2-naphthoic acid;
- 2-hydroxy-1-naphthoic acid;
- 3-hydroxy-2-pyridinecarboxylic acid;
- 4-hydroxy-7-methyl-3-naphthyridinecarboxylic acid;
- 2-hydroxy-3-pyridinecarboxylic acid;
- 2-hydroxybenzoic acid;
- 2,5-dihydroxybenzoic acid; and
- 2-hydroxy-5-phenylbenzoic acid (see E below)

there were obtained the following compounds:

- Compound 2 of Tables I and II;
- Compound 9 of Tables I and II;
- Compound 14 of Tables I and II;
- 5 Compound 15 of Tables I and II;
- Compound 16 of Tables I and II;
- Compound 17 of Tables I and II;
- Compound 18 of Tables I and II;
- Compound 19 of Tables I and II;
- 10 Compound 20 of Tables I and II; and
- Compound 21 of Tables I and II.

C. Preparation of Other Compounds of Formula I

By following the procedure of Part A above and  
15 substituting Fmoc-D-proline for Fmoc-L-proline,  
there was obtained Compound 12 of Tables III.

D. Preparation of Additional Compounds of Formula I

Compound 13 of Tables I and II was prepared by first  
20 coupling Fmoc-L-phenylalanine to the resin in an analogous  
procedure to that used to couple Fmoc-L-proline in part A  
above. The Fmoc group was then removed and the phenylalanine  
was coupled to Fmoc-L-proline. The remainder of the synthesis  
of Compound 13 was performed using the techniques described in  
25 Part A above.

Similarly, the preparation of Compounds 6 and 7 of  
Tables I and II involved coupling Fmoc-piperidine-3-carboxylic  
acid or Fmoc-piperidine-4-carboxylic acid, respectively, to the  
resin, removing the Fmoc group, and then coupling Fmoc-L-  
30 proline. The remainder of the syntheses were performed as  
described in Part A above.

E. Preparation 2-hydroxy-5-phenylbenzoic acid

2-hydroxy-5-phenylbenzoic acid was synthesized as  
35 follows. A solution of tetrahydropyran (THP)-protected 4-  
phenylphenol (474 mg. 2 mmol., prepared from 4-phenylphenol by  
standard methods described in Green and Wuts at pp. 31-34) in  
anhydrous THF (5 mL) was cooled to -78°C and treated with n-



Butyllithium (2 mL of 1.6 M in hexanes). The reaction mixture was stirred and warmed to ambient temperature during which a tan suspension formed. After 2 hours, the reaction mixture was cooled to  $-78^{\circ}\text{C}$  and treated with excess anhydrous  $\text{CO}_2$  for several minutes. The reaction mixture was stirred and warmed to ambient temperature. After 2 hours, the reaction mixture was partitioned between ethyl ether and aqueous 1N NaOH. The aqueous layer was cooled to  $0^{\circ}\text{C}$  with ice and acidified to pH 2 with aqueous 1N HCl. The aqueous solution was partitioned with methylene chloride. The organic layer was dried over magnesium sulfate. Concentration *in vacuo* yielded crude product as an off-white solid (258 mg, 2-hydroxy-5-phenylbenzoic acid). NMR spectroscopy of the crude material was consistent with the expected structure.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  8.04 (d,  $J=3\text{Hz}$ , 1H), 7.76 (dd,  $J=9$ , 3Hz, 1H), 7.62 (d,  $J=7\text{Hz}$ , 2H), 7.44 (t,  $J=7\text{Hz}$ , 2H), 7.32 (t,  $J=7\text{Hz}$ , 1H), 7.00 (d,  $J=9\text{Hz}$ , 1H). Mass calculated: 466.5. Mass found 467.2. LC Gradient 25-45% acetonitrile over 13 minutes. LC retention time 3.5 minutes. The material was used without further purification.

20

### III. Solution Phase Synthesis of Compounds of Formula I

#### A. Preparation of Compound 10 of Tables I and III

To 20 g of (L)-phenylalanine cooled to  $-5^{\circ}\text{C}$  in a salt/ice bath was added 74 mL concentrated sulfuric acid. The temperature of the reaction mixture was allowed to equilibrate and then 9.4 mL of concentrated nitric acid was added dropwise over five minutes. The reaction mixture was stirred for 30 minutes and was then added to 700 mL of crushed ice/water. The pH was adjusted to 8-9 with concentrated ammonium hydroxide. The solution was allowed to crystallize at room temperature and was then cooled overnight at  $4^{\circ}\text{C}$ . The light yellow crystalline product was collected over hardened filter paper, washed with cold water, and dried. The product was recrystallized in hot water and the filtrate was evaporated and recrystallized (2x) with an overall yield of 55%. MP:  $218-222^{\circ}\text{C}$  (crude). TLC:  $R_f$  (F) 0.18. NMR: ( $\text{D}_2\text{O}/\text{NaOD}$ ),  $\gamma$  2.99 (dd,  $J=13.4$ , 7.2 Hz, 1 H), 3.10 (dd,  $J=13.4$ , 6.0 Hz, 1 H), 3.57 (dd,  $J=7.2$ , 6.0 Hz, 1 H), 7.48 (d,  $J=8.6$  Hz, 2 H), 8.21 (d,  $J=8.7$  Hz, 2 H).

t-Butyloxycarbonyl (BOC)-protected amino acids can be prepared using standard techniques. See, e.g., Greene and Wuts Protective Groups in Organic Synthesis, 2nd Ed., John Wiley & Sons, Inc.: New York, pp. 327-328 (1991).

5 To a -20°C solution of BOC-protected p-nitrophenylalanine (3.00 g, 9.67 mmol) in anhydrous methylene chloride (5 mL) was added N-methylmorpholine (NMM, 1.07 mL, 9.67 mmol), followed by isobutyl chloroformate (1.26 mL, 9.67 mmol). The reaction mixture was stirred for 15 minutes at  
10 -20°C. To the mixture was then added methyl L-proline hydrochloride (L-Pro-OMe, 1.60 g, 9.67 mmol) as a solid, followed by additional NMM (1.07 mL, 9.67 mmol). After stirring for one hour at -20°C and one hour at room temperature, the reaction mixture was concentrated *in vacuo* and  
15 was then diluted with ethyl acetate. The ethyl acetate solution was washed with 10% aqueous citric acid, water, and brine, dried over magnesium sulfate, and concentrated *in vacuo*. Column chromatography yielded 2.4 g of product (>60% yield).

The BOC group was removed by treatment with HCl in  
20 dioxane using standard conditions. See, e.g., Greene and Wuts, *id* at 328-329. To a solution of 1-hydroxy-2-naphthoic acid (160 mg, 0.84 mmol) in methylene chloride and N,N-dimethylformamide (1:1, 3 mL) was added 1-hydroxybenzotriazole (159 mg, 1.17 mmol). The solution was cooled to 0°C and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI,  
25 161 mg, 0.84 mmol) was added. After the solution was stirred for 45 minutes at 0°C, a solution of the dipeptide (300 mg, 0.84 mmol) prepared above was added, followed by NMM (93 µl, 0.84 mmol). The reaction mixture was stirred for one hour at  
30 0°C and 45 minutes at room temperature. The reaction mixture was concentrated *in vacuo* and diluted with ethyl acetate and water. The organic layer was separated, washed (4x) with water and then brine, dried, and concentrated *in vacuo*. Chromatography yielded the coupled product (300 mg, 73%).

35 To a solution of the coupled product (410 mg, 0.83 mmol) in ethyl acetate (20 mL) was added glacial acetic acid (10 drops) and 10% palladium on activated carbon (100 mg). This solution was hydrogenated for five hours. The reaction

mixture was then filtered over celite and the solution was concentrated *in vacuo* to yield 375 mg (98%) of product which was used without further purification.

To a solution of the product of the above  
5 hydrogenation reaction (200 mg, 0.45 mmol) in anhydrous methanol (10 mL) was added formamidinesulfonic acid (118 mg, 0.95 mmol, prepared by oxidation of commercially available formamidinesulfinic acid following the procedure outlined in Maryanoff et al. (1986) J. Org. Chem. 1882). The reaction  
10 mixture was stirred for three days. The reaction mixture was concentrated *in vacuo* and the product was isolated by chromatography (eluting with methanol:methylene chloride) to yield 17 mg of desired product, Compound 10 of Table III.

15 IV. Alternate Solution Phase Synthesis of Compounds of Formula I

A. Preparation of Compound 3 of Tables I and II

A solution of 1-benzyloxy-2-naphthoic acid (2.247 g)  
20 in thionyl chloride (15 mL) was refluxed for four hours. Excess reagent was evaporated *in vacuo*. The residue was diluted with anhydrous DMF (9 mL) and concentrated to about 2 mL. To this solution was added 4-dimethylaminopyridine (1.02 g). The mixture was stirred at ambient temperature over  
25 night. The resulting pyridinium complex was added to a stirred solution of nitro-L-arginine (1.77 g), tetramethylguanidine (1 mL) and lithium chloride (4.56 g) in DMF (60 mL). The mixture was stirred at ambient temperature for 48 hours. Most of the DMF was evaporated *in vacuo* and the residue was  
30 partitioned between 1.5 normal (N) aqueous HCl and ethyl acetate. The combined organic layers were washed with saturated aqueous sodium chloride solution and dried over magnesium sulfate. Concentration *in vacuo* yielded crude product (3.5 g, 1-benzyloxy-2-naphthoyl-N-nitro-L-arginine acid). TLC:  
35  $R_f$  of product is 0.22 (using silica gel plate and eluting with 10% methanol/acetic acid in chloroform). This material was used without further purification.

A solution of 1-benzyloxy-2-naphthoyl-N-nitro-L-arginine acid (150 mg) in anhydrous DMF (5 mL) was cooled to

-25°C and treated with isobutyl chloroformate (0.053 mL). The reaction mixture was stirred and warmed to room temperature. After 20 minutes, solid L-proline amide (as its HCl salt, 65.9 mg) was added. The mixture was stirred for 16 hours and  
5 was then diluted with ethyl acetate. The ethyl acetate solution was washed with 5% aqueous citric acid, saturated aqueous sodium bicarbonate, and saturated aqueous sodium chloride and dried over magnesium sulfate. The solution was concentrated *in vacuo* and the residue was diluted in ethanol  
10 (40 mL, containing 10% methanol and 10% acetic acid) and hydrogenated at 52 pounds per square inch (psi) over 10% palladium on carbon (10 mg) until the reaction was complete (TLC on silica gel, eluting with chloroform:methanol 9:1) as indicated by disappearance of the faster moving spot. The  
15 reaction mixture was filtered through celite, washed with ethanol and methanol. The mixture was concentrated *in vacuo* and purified using high-pressure liquid chromatography to yield 1-hydroxy-2-naphthoyl-L-arginyl-L-proline amide (41 mg, electrospray mass spectrum:  $M^{+1} = 441$ ), isolated as its  
20 trifluoroacetate salt.

B. Preparation of Compound 1 of Table III

By following the procedure of Part A above and substituting 2-benzyloxy-1-naphthoic acid for 1-benzyloxy-2-  
25 naphthoic acid and sarcosine amide for L-proline amide, there was obtained Compound 1 of Table III.

C. Preparation of Other Compounds of Formula I

By following the procedures of Part B above and  
30 substituting for sarcosine amide the following compounds:

L-proline-N,N-dimethylamide;

L-proline methyl ester; and

trans-4-hydroxy-L-proline methyl ester;

there were obtained the following compounds:

35 Compound 5 of Tables I and II;

Compound 8 of Tables I and II; and

Compound 4 of Tables III.

## V. Physical Data

Table I

	Compound	Mass Calculated	Mass Found ( $M^{+1}$ ) <sup>1</sup>	LC Gradient <sup>2</sup>	LC Retention Time (min)
5	1	414.24	415.1	D	5.8
	2	404.26	405.5	B	11.5
	3	440.26	441.1	A	10.3
	4	471.26	472.2	D	6.2
10	5	468.29	469.1	D	6.8
	6 <sup>3</sup>	551.34	552.3	D	7.5
	7	551.34	552.1	D	7.3
	8	455.26	456.0	D	7.0
	9	442.26	443.0	D	5.5
15	10	-	-	-	-
	11	456.26	456.7	C	24.4
	12	440.26	441.2	C	25.4
	13	587.34	588.0	C	26.9
	14	440.26	440.5	A	9.0
20	15	440.26	440.2 <sup>4</sup>	B	8.4
	16	391.5	391.8 <sup>4</sup>	D	5.6
	17	456.49	456.7 <sup>4</sup>	B	9.0
	18	393.22	391.8 <sup>4</sup>	B	5.5
	19	390.4	391.1	A	4.7
25	20	406.4	407.2	C	20.7
	21	466.5	467.2	E	3.5

1. All masses were determined using a Finnigan electrospray mass spectrometer unless otherwise noted.

30

2. A binary gradient system consisting of acetonitrile and water was used in all cases. All eluents contain 0.1% trifluoroacetic acid. A: 20-45% acetonitrile over 13 minutes. B: 10-35% acetonitrile over 13 minutes. C: 5-95% acetonitrile over 45 minutes. D: 2-95% acetonitrile over 10 minutes. E: 25-45% acetonitrile over 13 minutes.

35

3. The material tested was a mixture of diastereomers.

4. Masses determined using a Biolon mass spectrometer. Values represent  $M^+$ .

5

#### VI. In Vitro Tryptase Inhibition Assay

Compounds (approx. 1 mg) to be assayed against tryptase were reconstituted in dimethylsulfoxide (DMSO) and diluted 1:10 into buffer containing 50 millimolar (mM) Tris-HCl at pH 8.2, and containing 100 mM NaCl, 0.05% Tween-20. Seven additional 3-fold dilutions were made from the initial dilution into the same buffer supplemented with 10% DMSO. Aliquots (50  $\mu$ l) from each of the eight dilutions in the series were transferred to individual wells in a 96-well U-bottom microtiter plate. Tryptase (25  $\mu$ l; 0.5 nM final concentration) was added to each well and the samples were mixed and incubated for 1 hr at room temperature under either ambient light (i.e., the light available in the room during the experiment) or under controlled "light" or "dark" conditions. "Light conditions" refers herein to a light intensity of 400-450 footcandles from a fluorescent bulb. "Dark" refers herein to sample containers wrapped in aluminum foil. The enzyme reaction was then initiated with the addition of the synthetic tripeptide substrate, tosyl-GlyProLys-p-nitroanilide (25  $\mu$ l; 0.5 mM final concentration). The microtiter plates were immediately transferred to a UV/MAX Kinetic Microplate Reader (Molecular Devices) and hydrolysis of the chromogenic substrate was followed spectrophotometrically at 405 nM for 5 minutes. The enzyme assays routinely yielded linear progress curves under these conditions. Initial velocity measurements, calculated from progress curves by a kinetic analysis program called "BatchKi" (available commercially from Biokin Ltd., of Madison WI), were used to determine apparent inhibition constants for each inhibitor. This program is designed to perform regressions and curve-fitting of non-linear data.










Table II lists the inhibition constants ( $K_i$ ', micromolar ( $\mu$ M)) which were determined for several of the compounds of the present invention, wherein  $R^1$  is hydrogen;  $R^2$  is hydrogen;  $R^3$  is  $-(CH_2)_3-NH-(C=NH)-NH_2$ ;  $R^4$  and  $R^5$ , together with the nitrogen and carbon to which they are attached form a

40







five-membered heterocycle; and  $R^6$  is hydrogen. According to the present invention, a compound was termed "active" or effective as a tryptase inhibitor when its  $K_i'$  was less than 1000  $\mu M$ . Unlike  $K_i$ ,  $K_i'$  may not be a true dissociation constant of the enzyme-inhibitor complex;  $K_i'$  is equal to  $K_i$  for a noncompetitive inhibitor and is directly proportional to  $K_i$  for competitive and uncompetitive inhibitors.

Exposure of solutions of the test compounds in the assay medium to light may decrease the  $K_i'$ . Generally, the assay conditions are light sensitive and variations in ambient light intensity may affect assay reproducibility. As a result of this potential source of error, some assays were run under dark conditions. The  $K_i$ 's reported in Tables II and III without parentheses were determined under conditions of ambient light.  $K_i$ 's reported with parentheses were determined under "dark" conditions as defined above.  $K_i$ 's reported with an asterisk (\*) were measured under "light" conditions as defined above (i.e., at a light intensity of 400-450 footcandles).

Table II

Compound	$K_i'$	$R^7$	Ar
2	>500	$-\text{NH}_2$	
3	0.3* (82)	$-\text{NH}_2$	
5	15	$-\text{N}(\text{CH}_3)_2$	
6	82		
7	664		
8	208	$-\text{OCH}_3$	
9	23* (56)	$-\text{NH}_2$	



Comp und	$K_i'$	$R^7$	Ar
13	0.6* (110)		
14	752* (707)	-NH <sub>2</sub>	
15	0.84	-NH <sub>2</sub>	
16	8* (63)	-NH <sub>2</sub>	
17	900	-NH <sub>2</sub>	

5

10









Compound	$K_i'$	$R^7$	Ar
18	452	$-NH_2$	
19	8* (2450)	$-NH_2$	
20	20	$-NH_2$	
21	18* (149)	$-NH_2$	

Table III lists the inhibition constants ( $K_i'$ , micromolar ( $\mu M$ )) which were determined for several of the compounds of the present invention. According to the present invention, a compound was termed "active" or effective as a tryptase inhibitor when its  $K_i'$  was less than 1000  $\mu M$ .


Table III

	Compound	K <sub>i</sub> '	Structure
5  10	1	3* (637)	
15  20	4	749	

5

C mpound	K <sub>i</sub> ' (nM)	Structure
10	2.5* (18)	
11	0.3* (40)	

10

Comp und	$K_i'$	Structure
12	2.5* (818)	

VII. Additional Characterization of Tryptase Inhibition by Compound 3 of Tables I and II

A. pH Dependency

Compound 3 was assayed against tryptase at four different pH's in the following buffer system: 120 mM NaCl, 2.7 mM KCl, 0.13 mM  $\text{NaH}_2\text{PO}_4$ , 0.896 mM  $\text{Na}_2\text{HPO}_4$ . The final buffer system used included 0.05% of Tween-20. Table IV shows  $K_i'$  ( $\mu\text{M}$ ) for Compound 3 at different pH's.

Table IV

pH	<u>6.5</u>	<u>7.0</u>	<u>7.5</u>	<u>8.0</u>
$K_i'$ ( $\mu\text{M}$ )	14	8.0	5.4	6.1

B. Optimized Assay at pH 7.5

The inventors have discovered that Compound 3 is less sensitive to light in the following buffer system: 120 mM NaCl, 2.7 mM KCl, 0.13 mM  $\text{NaH}_2\text{PO}_4$ , 0.896 mM  $\text{Na}_2\text{HPO}_4$  and 0.05% Tween-20 at a pH of 7.5. Thus, the assay procedure described above was performed wherein the dilution of Compound 3 and the addition of tryptase was done under ambient light at room temperature (22-26°C). The assay plate was wrapped in aluminum

foil for a 1-hr incubation before substrate addition, which was also done under ambient light. The steps of dilution and the addition of tryptase must be completed within 5 minutes. In addition, Compound 3 must be reconstituted in DMSO.

5

#### VIII. In Vivo Testing

The allergic sheep model of asthma was employed in these studies. These methods have been published previously (see Abraham et al. (1983) Am. Rev. Respir. Dis. 128:839-844; Allegra et al. (1983) J. Appl. Physiol. 55:726-730; Russi et al. (1985) J. Appl. Physiol. 59:1416-1422; Soler et al. ((1989) J. Appl. Physiol. 67:406-413. Each sheep served as its own control. Body weights for these animals ranged from 20-50 kilograms.

15 In these studies, 9 mg of Compound 3 of Tables I and II was dissolved in 3 mL buffered saline and the total solution delivered as an aerosol 0.5 hours before, 4 hours after, and 24 hours after antigen challenge (total dose = 27; n= 6). Compound 3 exhibited a tendency to reduce the early response, and attenuated the late response as shown in Figure 1. In the control (vehicle only) trial, antigen challenge caused peak early and late increases above baseline in specific lung resistance (SRL) of  $374 \pm 104\%$  and  $212 \pm 21\%$  (mean  $\pm$  SE). In contrast, when the sheep were treated with Compound 3 of Tables I and II, the early and late increases in SRL were  $280 \pm 39\%$  and  $72 \pm 9\%$  ( $p < 0.05$  vs. control, Wilcoxon signed rank test). The peak early response was taken as the average of the maximum values occurring immediately post-challenge. Peak late responses were calculated by averaging the maximum response values obtained for each animal within the 6 - 8 hour time period. This approach is conservative and eliminates the possible reduction in the late response due simply to averaging.

35 Twenty-four hours after antigen challenge in both the control and drug trial, the sheep developed airway hyperresponsiveness. (Airway hyperresponsiveness is expressed as PC400, i.e., the concentration of carbachol that causes a 400% increase in SRL. Thus a decrease in the PC400 indicates


that the airways have become hyperresponsive.) Compound 3 of Tables I and II blocked the 24 hour hyperresponsiveness. See Figure 2. Baseline PC400 was  $22.0 \pm 3.7$  breath units in the control trial, falling to  $9.1 \pm 1.3$  breath units after antigen challenge (p < 0.05 vs. baseline). In contrast, in the drug trial, PC400 was unchanged relative to baseline ( $16.0 \pm 3.8$  breath units vs.  $17.6 \pm 3.5$  breath units after antigen challenge). Thus treatment with Compound 3 of Tables I and II resulted in a statistically significant improvement in airway function in allergen challenged sheep.

The disclosures in this application of all articles and references, including patents, are incorporated herein by reference.

It is to be understood that the above description is intended to be illustrative and not restrictive. Many embodiments will be apparent to those of skill in the art upon reviewing the above description. The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with reference to the appended claims, along with the full scope of equivalents to which such claims are entitled.

WHAT IS CLAIMED IS:


1. A compound having the formula:




- or a pharmaceutically acceptable salt thereof, wherein:  
Ar is hydroxyl substituted aryl or hydroxyl substituted heteroaryl, wherein the hydroxyl is positioned ortho to the amide side chain and wherein if Ar is hydroxyl substituted aryl, the aromatic ring bearing the amide side chain is not substituted with halogen and does not bear a lower alkyl group on the position ortho to the hydroxyl;  
R<sup>1</sup> is hydrogen, lower alkyl, arylalkyl, or heteroarylalkyl;  
R<sup>2</sup> is hydrogen or lower alkyl;  
R<sup>3</sup> is selected from the group consisting of:




47




5




10




15




20




25



30



and



wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; r is an integer from 0-5; s is an integer from 0-2; t is an integer from 1-3; u is 1 or 2; v is an integer from 3-6; and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is  $-\text{NH}-$  or  $-\text{CH}_2-$ ;

R<sup>4</sup> is lower alkyl, substituted arylalkyl, or substituted heteroarylalkyl, R<sup>5</sup> and R<sup>6</sup> are independently selected from the group consisting of hydrogen, lower alkyl, substituted arylalkyl, and substituted heteroarylalkyl; or R<sup>4</sup> and R<sup>5</sup> together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and R<sup>6</sup> is hydrogen; or R<sup>4</sup> and R<sup>6</sup> together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and R<sup>5</sup> is hydrogen; and

R<sup>7</sup> is -OR<sup>8</sup> or -NR<sup>8</sup>R<sup>9</sup>, wherein R<sup>8</sup> and R<sup>9</sup> are independently selected from the group consisting of hydrogen, lower alkyl, aryl, arylalkyl, or heteroarylalkyl or R<sup>8</sup> and R<sup>9</sup> together with the nitrogen to which they are attached, form a substituted 5-membered or 6-membered heterocycle.

2. The compound of Claim 1 wherein:


Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;

R<sup>1</sup> is hydrogen;


R<sup>2</sup> is hydrogen;

R<sup>3</sup> is selected from the group consisting of:


49




5



10



15



wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is  $-\text{NH}-$  or  $-\text{CH}_2-$ ;

$\text{R}^4$  is lower alkyl and  $\text{R}^5$  and  $\text{R}^6$  is hydrogen; or  $\text{R}^4$  and  $\text{R}^5$  together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $\text{R}^6$  is hydrogen;

and  $\text{R}^7$  is  $-\text{OH}$ ,  $-\text{OCH}_3$ ,  $-\text{NH}_2$ ,  $-3'\text{-aminocarboxy-1'-piperidyl}$ , or  $-\text{N}(\text{CH}_3)_2$ .

3. The compound of Claim 2 wherein Ar is 1-hydroxy-2-naphthyl,  $\text{R}^1$  is hydrogen,  $\text{R}^2$  is hydrogen,  $\text{R}^3$  is  $-(\text{CH}_2)_3\text{NH}(\text{CNH})\text{NH}_2$ ,  $\text{R}^4$  and  $\text{R}^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $\text{R}^6$  is hydrogen, and  $\text{R}^7$  is  $-\text{NH}_2$ .

4. The compound of Claim 2 wherein Ar is 2-hydroxy-1-naphthyl,  $\text{R}^1$  is hydrogen,  $\text{R}^2$  is hydrogen,  $\text{R}^3$  is

$-(CH_2)_3NH(CNH)NH_2$ ,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is  $-NH_2$ .

5                    5. An aerosol composition for the treatment of immunomediated inflammatory disorders comprising a compound of Claim 1 in an aerosolized pharmaceutically acceptable carrier solution or dry powder.


10                   6. The composition of Claim 5 wherein:

Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;


$R^1$  is hydrogen;

$R^2$  is hydrogen;


15                    $R^3$  is selected from the group consisting of:




20



25



30



35                   wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is  $-\text{NH}-$  or  $-\text{CH}_2-$ ;

$R^4$  is lower alkyl and  $R^5$  and  $R^6$  is hydrogen; or  $R^4$  and  $R^5$  together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $R^6$  is hydrogen;

5                   and  $R^7$  is -OH, -OCH<sub>3</sub>, -NH<sub>2</sub>, -3'aminocarboxy-1'-piperidyl, or -N(CH<sub>3</sub>)<sub>2</sub>.

7.     The composition of Claim 6 wherein Ar is 1-hydroxy-2-naphthyl,  $R^1$  is hydrogen,  $R^2$  is hydrogen,  $R^3$  is  
10    -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is -NH<sub>2</sub>.

8.     The composition of Claim 6 wherein Ar is 2-hydroxy-1-naphthyl,  $R^1$  is hydrogen,  $R^2$  is hydrogen,  $R^3$  is  
15    -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is -NH<sub>2</sub>.

9.     The composition of Claim 5 wherein said inflammatory disorder of the respiratory tract is asthma.

10.    The composition of Claim 5 wherein said inflammatory disorder of the respiratory tract is allergic  
25    rhinitis.

11.    The composition of Claim 5 wherein said compound of Claim 1 is present in said carrier solution in a concentration of from 0.1 to 30 mg/mL.

30

12.    The composition of Claim 5 further comprising a  $\beta$ -adrenergic agonist compound.

13.    The composition of Claim 12 wherein said  $\beta$ -adrenergic agonist compound is selected from the group consisting of albuterol, terbutaline, formoterol, fenoterol, and prenaline.

35

14. The composition of Claim 5 further comprising an antiinflammatory corticosteroid.

5 15. The composition of Claim 14 wherein said antiinflammatory corticosteroid is selected from the group consisting of beclomethasone, triamcinolone, flurisolide, and dexamethasone.

10 16. The composition of Claim 5 further comprising ipratropium bromide.

15 17. A pharmaceutical composition comprising a compound of Claim 1 in combination with a pharmaceutically acceptable carrier.

18. The composition of Claim 17 wherein the pharmaceutically acceptable carrier comprises a non-toxic, pharmaceutically acceptable topical carrier.

20 19. The composition of Claim 17 wherein:

Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;


R<sup>1</sup> is hydrogen;

R<sup>2</sup> is hydrogen;


25 R<sup>3</sup> is selected from the group consisting of:

30


35




5



10



15



wherein m is an integer from 3-6, n is an integer  
 20 from 0-3, p is an integer from 0-2, q is an integer from 0-2;  
 and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is -  
 NH- or  
 $-\text{CH}_2-$ ;

$\text{R}^4$  is lower alkyl and  $\text{R}^5$  and  $\text{R}^6$  is hydrogen; or  $\text{R}^4$  and  
 25  $\text{R}^5$  together with the nitrogen and carbon to which they are  
 attached form a substituted 4-membered, 5-membered, or  
 6-membered heterocycle and  $\text{R}^6$  is hydrogen;

and  $\text{R}^7$  is  $-\text{OH}$ ,  $-\text{OCH}_3$ ,  $-\text{NH}_2$ , -3'aminocarboxy-1'-  
 piperidyl, or  $-\text{N}(\text{CH}_3)_2$ .

30




20. The composition of Claim 19 wherein Ar is 1-  
 hydroxy-2-naphthyl,  $\text{R}^1$  is hydrogen,  $\text{R}^2$  is hydrogen,  $\text{R}^3$  is  
 $-(\text{CH}_2)_3\text{NH}(\text{CNH})\text{NH}_2$ ,  $\text{R}^4$  and  $\text{R}^5$  along with the nitrogen and carbon  
 to which they are attached form a substituted 5-membered  
 35 heterocycle,  $\text{R}^6$  is hydrogen, and  $\text{R}^7$  is  $-\text{NH}_2$ .

21. The composition of Claim 19 wherein Ar is 2-  
 hydroxy-1-naphthyl,  $\text{R}^1$  is hydrogen,  $\text{R}^2$  is hydrogen,  $\text{R}^3$  is

$-(CH_2)_3NH(CNH)NH_2$ ,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is  $-NH_2$ .

- 5                    22. An aerosol device comprising:  
                       a compound of Claim 1 in a pharmaceutically  
                       acceptable carrier solution or dry powder, and  
                       means for converting said solution or dry powder into  
 10                   an aerosol form suitable for inhalation.

- 10                   23. The device of Claim 22 wherein:  
                       Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl,  
                       3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;  
                        $R^1$  is hydrogen;  
 15                    $R^2$  is hydrogen;  
                        $R^3$  is selected from the group consisting of:



- 35                   wherein m is an integer from 3-6, n is an integer  
                       from 0-3, p is an integer from 0-2, q is an integer from 0-2;  
                       and w is an integer from 0-3; A is  $-CH=CH-$  or  $-C \equiv C-$ ; and X is  $-$   
                        $NH-$  or  $-CH_2-$ ;



$R^4$  is lower alkyl and  $R^5$  and  $R^6$  is hydrogen; or  $R^4$  and  $R^5$  together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $R^6$  is hydrogen;

5                   and  $R^7$  is -OH, -OCH<sub>3</sub>, -NH<sub>2</sub>, -3'aminocarboxy-1'-piperidyl, or -N(CH<sub>3</sub>)<sub>2</sub>.

24. The device of Claim 23 wherein Ar is 1-hydroxy-2-naphthyl,  $R^1$  is hydrogen,  $R^2$  is hydrogen,  $R^3$  is  
10   -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is -NH<sub>2</sub>.

25. The device of Claim 23 wherein Ar is 2-hydroxy-1-naphthyl,  $R^1$  is hydrogen,  $R^2$  is hydrogen,  $R^3$  is  
15   -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is -NH<sub>2</sub>.

20                   26. The device of Claim 22 wherein said compound is present in said carrier solution in a concentration of from 0.1 to 30 mg/mL.

25                   27. The device of Claim 22 further comprising a  $\beta$ -adrenergic agonist compound.

28. The device of Claim 27 wherein said  $\beta$ -adrenergic agonist compound is selected from the group consisting of albuterol, terbutaline, formoterol, fenoterol, and prenaline.  
30

29. The device of Claim 22 further comprising an antiinflammatory corticosteroid.

30. The device of Claim 29 wherein said  
35   antiinflammatory corticosteroid is selected from the group consisting of beclomethasone, triamcinolone, flurisolide, and dexamethasone.

31. The device of Claim 22 further comprising ipratropium bromide.

32. A device for the treatment of allergic rhinitis comprising:

a compound of Claim 1 in a pharmaceutically acceptable carrier solution, and

means for converting said solution into an aerosol form suitable for intranasal administration.

10

33. The device of Claim 32 wherein:

Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;


R<sup>1</sup> is hydrogen;

15


R<sup>2</sup> is hydrogen;

R<sup>3</sup> is selected from the group consisting of:



20



25



30



35

wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; and w is an integer from 0-3; A is -CH=CH- or -C≡C-; and X is -NH- or -CH<sub>2</sub>-;

$R^4$  is lower alkyl and  $R^5$  and  $R^6$  is hydrogen; or  $R^4$  and  $R^5$  together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $R^6$  is hydrogen;

5                   and  $R^7$  is -OH, -OCH<sub>3</sub>, -NH<sub>2</sub>, -3'aminocarboxy-1'-piperidyl, or -N(CH<sub>3</sub>)<sub>2</sub>.

34. The device of Claim 33 wherein Ar is 1-hydroxy-2-naphthyl,  $R^1$  is hydrogen,  $R^2$  is hydrogen,  $R^3$  is  
10   -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is -NH<sub>2</sub>.

35. The device of Claim 33 wherein Ar is 2-hydroxy-1-naphthyl,  $R^1$  is hydrogen,  $R^2$  is hydrogen,  $R^3$  is  
15   -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is -NH<sub>2</sub>.

20                   36. A method for treating an immunomediated inflammatory disorder of the respiratory tract, said method comprising administering to a mammal an inhalant composition comprising a compound of Claim 1 in an aerosolized pharmaceutically acceptable carrier solution or dry powder.

25

37. The method of Claim 36 wherein:  
Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;


$R^1$  is hydrogen;

30                    $R^2$  is hydrogen;


$R^3$  is selected from the group consisting of:

35


58




5



10



15



wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; and w is an integer from 0-3; A is -CH=CH- or -C≡C-; and X is -NH- or -CH<sub>2</sub>-;

R<sup>4</sup> is lower alkyl and R<sup>5</sup> and R<sup>6</sup> is hydrogen; or R<sup>4</sup> and R<sup>5</sup> together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and R<sup>6</sup> is hydrogen;

and R<sup>7</sup> is -OH, -OCH<sub>3</sub>, -NH<sub>2</sub>, -3'aminocarboxy-1'-piperidyl, or -N(CH<sub>3</sub>)<sub>2</sub>.

38. The method of Claim 37 wherein Ar is 1-hydroxy-2-naphthyl, R<sup>1</sup> is hydrogen, R<sup>2</sup> is hydrogen, R<sup>3</sup> is -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>, R<sup>4</sup> and R<sup>5</sup> along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle, R<sup>6</sup> is hydrogen, and R<sup>7</sup> is -NH<sub>2</sub>.

39. The method of Claim 37 wherein Ar is 2-hydroxy-1-naphthyl, R<sup>1</sup> is hydrogen, R<sup>2</sup> is hydrogen, R<sup>3</sup> is

$-(\text{CH}_2)_3\text{NH}(\text{CNH})\text{NH}_2$ ,  $\text{R}^4$  and  $\text{R}^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $\text{R}^6$  is hydrogen, and  $\text{R}^7$  is  $-\text{NH}_2$ .

5           40. The method of Claim 36 wherein said compound is present in said carrier solution in a concentration of from 0.1 to 30 mg/mL.

10           41. The method of Claim 36 further comprising a  $\beta$ -adrenergic agonist compound.

15           42. The method of Claim 41 wherein said  $\beta$ -adrenergic agonist compound is selected from the group consisting of albuterol, terbutaline, formoterol, fenoterol, and prenaline.

          43. The method of Claim 36 further comprising an antiinflammatory corticosteroid.

20           44. The method of Claim 43 wherein said antiinflammatory corticosteroid is selected from the group consisting of beclomethasone, triamcinolone, flurisolide, and dexamethasone.

25           45. The method of Claim 36 further comprising ipratropium bromide.

30           46. A method for treating immunomediated inflammatory skin conditions, said method comprising administering topically to a mammal a composition comprising a therapeutically effective amount of a compound of Claim 1 in a non-toxic, pharmaceutically acceptable topical carrier.

          47. The method of Claim 46 wherein:


35           Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;

$\text{R}^1$  is hydrogen;


$\text{R}^2$  is hydrogen;

$\text{R}^3$  is selected from the group consisting of:


60




5



10



15



wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; and w is an integer from 0-3; A is  $-\text{CH}=\text{CH}-$  or  $-\text{C}\equiv\text{C}-$ ; and X is  $-\text{NH}-$  or  $-\text{CH}_2-$ ;

$\text{R}^4$  is lower alkyl and  $\text{R}^5$  and  $\text{R}^6$  is hydrogen; or  $\text{R}^4$  and  $\text{R}^5$  together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and  $\text{R}^6$  is hydrogen;

and  $\text{R}^7$  is  $-\text{OH}$ ,  $-\text{OCH}_3$ ,  $-\text{NH}_2$ ,  $-3'$ aminocarboxy- $1'$ -piperidyl, or  $-\text{N}(\text{CH}_3)_2$ .

48. The method of Claim 47 wherein Ar is 1-hydroxy-2-naphthyl,  $\text{R}^1$  is hydrogen,  $\text{R}^2$  is hydrogen,  $\text{R}^3$  is  $-(\text{CH}_2)_3\text{NH}(\text{CNH})\text{NH}_2$ ,  $\text{R}^4$  and  $\text{R}^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $\text{R}^6$  is hydrogen, and  $\text{R}^7$  is  $-\text{NH}_2$ .

49. The method of Claim 47 wherein Ar is 2-hydroxy-1-naphthyl,  $\text{R}^1$  is hydrogen,  $\text{R}^2$  is hydrogen,  $\text{R}^3$  is

$-(CH_2)_3NH(CNH)NH_2$ ,  $R^4$  and  $R^5$  along with the nitrogen and carbon to which they are attached form a substituted 5-membered heterocycle,  $R^6$  is hydrogen, and  $R^7$  is  $-NH_2$ .

5                    50. A method for prophylactically treating an immunomediated inflammatory disorder of the respiratory tract, said method comprising administering to a mammal an inhalant composition comprising a prophylactically effective amount of a compound of Claim 1 in an aerosolized pharmaceutically  
10 acceptable carrier solution or dry powder.





51. The method of Claim 50 wherein:

Ar is 1-hydroxy-2-naphthyl, 2-hydroxyl-1-naphthyl, 3-hydroxy-2-pyridyl, or 2-hydroxy-3-quinoxalyl;

15                     $R^1$  is hydrogen;

$R^2$  is hydrogen;

$R^3$  is selected from the group consisting of:



35

wherein m is an integer from 3-6, n is an integer from 0-3, p is an integer from 0-2, q is an integer from 0-2; and w is an integer from 0-3; A is -CH=CH- or -C=C-; and X is -NH- or -CH<sub>2</sub>-;

- 5           R<sup>4</sup> is lower alkyl and R<sup>5</sup> and R<sup>6</sup> is hydrogen; or R<sup>4</sup> and R<sup>5</sup> together with the nitrogen and carbon to which they are attached form a substituted 4-membered, 5-membered, or 6-membered heterocycle and R<sup>6</sup> is hydrogen;  
and R<sup>7</sup> is -OH, -OCH<sub>3</sub>, -NH<sub>2</sub>, -3'aminocarboxy-1'-  
10 piperidyl, or -N(CH<sub>3</sub>)<sub>2</sub>.

52. The method of Claim 51 wherein Ar is 1-hydroxy-2-naphthyl, R<sup>1</sup> is hydrogen, R<sup>2</sup> is hydrogen, R<sup>3</sup> is -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>, R<sup>4</sup> and R<sup>5</sup> along with the nitrogen and carbon  
15 to which they are attached form a substituted 5-membered heterocycle, R<sup>6</sup> is hydrogen, and R<sup>7</sup> is -NH<sub>2</sub>.

53. The method of Claim 51 wherein Ar is 2-hydroxy-1-naphthyl, R<sup>1</sup> is hydrogen, R<sup>2</sup> is hydrogen, R<sup>3</sup> is -(CH<sub>2</sub>)<sub>3</sub>NH(CNH)NH<sub>2</sub>, R<sup>4</sup> and R<sup>5</sup> along with the nitrogen and carbon  
20 to which they are attached form a substituted 5-membered heterocycle, R<sup>6</sup> is hydrogen, and R<sup>7</sup> is -NH<sub>2</sub>.

54. The method of Claim 50 wherein said compound is  
25 present in said carrier solution in a concentration of from 0.1 to 30 mg/mL.

55. The method of Claim 50 further comprising a β-adrenergic agonist compound.  
30

56. The method of Claim 55 wherein said β-adrenergic agonist compound is selected from the group consisting of albuterol, terbutaline, formoterol, fenoterol, and prenaline.

- 35           57. The method of Claim 50 further comprising an antiinflammatory corticosteroid.



58. The method of Claim 57 wherein said antiinflammatory corticosteroid is selected from the group consisting of beclomethasone, triamcinolone, flurisolide, and dexamethasone.

5

59. The method of Claim 50 further comprising ipratropium bromide.

1/1




FIG. 1.




FIG. 2.

## INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/US 94/02706

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 5 C07K5/06 A61K37/02

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	L.B.SCHWARTZ ET AL. 'Monographs in Allergy' 1990, KARGER, BASEL *Neutral Proteases of Mast Cells* see page 51-66 ---	
A	US,A,4 201 863 (SHOSUKE ET AL.) 6 May 1980 ---	
A	US,A,4 456 595 (H.N.WELLER ET AL.) 26 June 1984 -----	

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

29 June 1994

Date of mailing of the international search report

03. 08. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Deffner, C-A

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/02706

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4201863	06-05-80	JP-C- 1167853	08-09-83
		JP-A- 51105046	17-09-76
		JP-B- 57047990	13-10-82
		JP-C- 1165631	26-08-83
		JP-A- 51105047	17-09-76
		JP-B- 57047991	13-10-82
		JP-C- 1264290	16-05-85
		JP-A- 52019653	15-02-77
		JP-B- 59041991	11-10-84
		JP-C- 1240709	26-11-84
		JP-A- 52031061	09-03-77
		JP-B- 59013500	30-03-84
		JP-C- 1240711	26-11-84
		JP-A- 52033662	14-03-77
		JP-B- 59013501	30-03-84
		JP-C- 1240714	26-11-84
		JP-A- 52042863	04-04-77
		JP-B- 59013503	30-03-84
		JP-C- 1189055	13-02-84
		JP-A- 52046059	12-04-77
		JP-B- 58022030	06-05-83
		JP-C- 1052091	30-06-81
		JP-A- 51125052	01-11-76
		JP-B- 55042981	04-11-80
		JP-C- 1052092	30-06-81
		JP-A- 51125053	01-11-76
		JP-B- 55042982	04-11-80
		JP-C- 1047480	28-05-81
		JP-A- 51125051	01-11-76
		JP-B- 55037992	01-10-80
		JP-C- 1067772	23-10-81
		JP-A- 51125260	01-11-76
		JP-B- 56008832	25-02-81
		US-A- 4258192	24-03-81
		CA-A- 1073914	18-03-80
		DE-A- 2550088	20-05-76
		FR-A, B 2290193	04-06-76
		GB-A- 1516668	05-07-78
		NL-A- 7512637	11-05-76
		SE-B- 431204	23-01-84

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/02706

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4201863		SE-A- 7512530	10-05-76
		US-A- 4055636	25-10-77
		US-A- 4055651	25-10-77
		US-A- 4041156	09-08-77
		US-A- 4046876	06-09-77
		US-A- 4070457	24-01-78
		US-A- 4062963	13-12-77
		US-A- 4066758	03-01-78
		US-A- 4049645	20-09-77
		US-A- 4069323	17-01-78
<hr/>			
US-A-4456595	26-06-84	NONE	
<hr/>			